

Mathematical
Inequalities
& **A**pplications

*With Compliments
of the Author*

Zagreb, Croatia

Volume 23, Number 4, October 2020

Chao-Ping Chen and Richard B. Paris

*An inequality involving the constant e and a
generalized Carleman-type inequality*

MATHEMATICAL INEQUALITIES & APPLICATIONS

AIMS AND SCOPE

Mathematical Inequalities & Applications (MIA, Math. Inequal. Appl.) brings together original research papers in all areas of mathematics, provided they are concerned with inequalities or their role. From time to time MIA will publish invited survey articles. Short notes with interesting results or open problems will also be accepted.

MIA is published quarterly, in January, April, July and October.

SUBMISSION

Manuscripts should be submitted electronically in PostScript or Adobe Acrobat PDF format to the MIA Editorial Office through MIA page: www.ele-math.com or by e-mail mia@ele-math.com

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the MIA regarding the article. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to MIA. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal, and that its submission for publication has been approved by all the authors.

PREPARATION OF MANUSCRIPT

Manuscripts should be written in English.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send final version of file(s) to the Editorial Office.

TITLE PAGE

A title page should include the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author and running head less than 60 characters. Authors should be listed alphabetically.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2000) with primary (and secondary) subject classification codes and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, grey scale only, preferably at resolution of 300 dpi or higher, sized to fit a journal page, and in EPS, TIFF or JPEG format. Figures should have a short descriptive title and should be numbered consecutively in order of appearance in the text. The position of each figure in the text should be indicated.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The author should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS

The author will receive proofs only once. Late return of the proofs will delay the article to a later issue.

PAGE CHARGES, OFFPRINTS

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from 'MIA' prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

MIA is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is not supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

Mathematical Inequalities & Applications
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
www.ele-math.com
e-mail: mia@ele-math.com
Fax: +385 1 6008799

AN INEQUALITY INVOLVING THE CONSTANT e AND A GENERALIZED CARLEMAN-TYPE INEQUALITY

CHAO-PING CHEN AND RICHARD B. PARIS

(Communicated by I. Perić)

Abstract. In this paper, we establish a double inequality involving the constant e . As an application, we give a generalized Carleman-type inequality.

1. Introduction

Let $a_n \geq 0$ for $n \in \mathbb{N} := \{1, 2, \dots\}$ and $0 < \sum_{n=1}^{\infty} a_n < \infty$. Then

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} a_n. \quad (1.1)$$

The constant e is the best possible. The inequality (1.1) was presented in 1922 in [3] by the Swedish mathematician Torsten Carleman and it is called Carleman's inequality. Carleman discovered this inequality during his important work on quasi-analytical functions.

Carleman's inequality (1.1) was generalized by Hardy [12] (see also [13, p. 256]) as follows: If $a_n \geq 0$, $\lambda_n > 0$, $\Lambda_n = \sum_{m=1}^n \lambda_m$ for $n \in \mathbb{N}$, and $0 < \sum_{n=1}^{\infty} \lambda_n a_n < \infty$, then

$$\sum_{n=1}^{\infty} \lambda_n (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < e \sum_{n=1}^{\infty} \lambda_n a_n. \quad (1.2)$$

Note that inequality (1.2) is usually referred to as a Carleman-type inequality or weighted Carleman-type inequality. In his original paper [12], Hardy himself said that it was Pólya who pointed out this inequality to him. For information about the history of Carleman-type inequalities, please refer to [15, 16, 18, 24].

In [4, 5, 6, 9, 10, 11, 14, 19, 20, 21, 22, 23, 26, 27, 28, 29, 30, 31], some strengthened and generalized results of (1.1) and (1.2) have been given by estimating the weight coefficient $(1 + 1/n)^n$. For example, Mortici and Jang [23] proved that for $0 < x \leq 1$,

$$\begin{aligned} e \left(1 - \frac{1}{2}x + \frac{11}{24}x^2 - \frac{7}{16}x^3 + \frac{2447}{5760}x^4 - \frac{959}{2304}x^5 \right) &< (1+x)^{1/x} \\ &< e \left(1 - \frac{1}{2}x + \frac{11}{24}x^2 - \frac{7}{16}x^3 + \frac{2447}{5760}x^4 \right). \end{aligned} \quad (1.3)$$

Mathematics subject classification (2010): 26D15.

Keywords and phrases: Carleman's inequality, weight coefficient.

This work was supported by Key Science Research Project in Universities of Henan (20B110007).

According to Pólya's proof of (1.1) in [25],

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} \leq \sum_{n=1}^{\infty} \left(1 + \frac{1}{n}\right)^n a_n, \quad (1.4)$$

and then the following strengthened Carleman's inequality can be derived directly from the right-hand side of (1.3):

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} \left(1 - \frac{1}{2n} + \frac{11}{24n^2} - \frac{7}{16n^3} + \frac{2447}{5760n^4}\right) a_n. \quad (1.5)$$

In this paper, we develop the double inequality (1.3) to produce a general result. As an application, we give a generalized Carleman-type inequality.

2. A double inequality involving the constant e

Brothers and Knox [2] (see also [17, 7]) derived, without a formula for the general term, the following expansion:

$$\left(1 + \frac{1}{x}\right)^x = e \left(1 - \frac{1}{2x} + \frac{11}{24x^2} - \frac{7}{16x^3} + \frac{2447}{5760x^4} - \frac{959}{2304x^5} + \frac{238043}{580608x^6} - \cdots\right) \quad (2.1)$$

for $x < -1$ or $x \geq 1$. Chen and Choi [7] gave an explicit formula for successively determining the coefficients. More precisely, these authors proved that

$$\left(1 + \frac{1}{x}\right)^x \sim e \sum_{j=0}^{\infty} (-1)^j b_j x^{-j} \quad (x \rightarrow \infty), \quad (2.2)$$

where the coefficients b_j are given by

$$b_0 = 1 \quad \text{and} \quad b_j = \sum_{k_1+2k_2+\cdots+jk_j=j} \frac{\left(\frac{1}{2}\right)^{k_1} \left(\frac{1}{3}\right)^{k_2} \cdots \left(\frac{1}{j+1}\right)^{k_j}}{k_1! k_2! \cdots k_j!} \quad (j \geq 1) \quad (2.3)$$

summed over all nonnegative integers k_j satisfying the equation $k_1 + 2k_2 + \cdots + jk_j = j$.

A recurrence relation for the coefficients b_j can be obtained by use of the result given in [8, Lemma 3]. This states that for a function $A(x)$ with asymptotic expansion $A(x) \sim \sum_{n=1}^{\infty} \alpha_n x^{-n}$ as $x \rightarrow \infty$, the composition $B(x) = \exp[A(x)]$ has the expansion $B(x) \sim \sum_{n=1}^{\infty} \beta_n x^{-n}$ as $x \rightarrow \infty$, where $\beta_0 = 1$ and

$$\beta_n = \frac{1}{n} \sum_{k=1}^n k \alpha_k \beta_{n-k} \quad (n \geq 1).$$

From the Maclaurin expansion

$$\frac{1}{x} \ln(1+x) = 1 + \sum_{j=1}^{\infty} \frac{(-1)^j x^j}{j+1} \quad (-1 < x \leq 1),$$

it therefore follows (upon replacing x by $1/x$) that the coefficients b_j in (2.2) are given by the recurrence relation

$$b_0 = 1 \quad \text{and} \quad b_j = \frac{1}{j} \sum_{k=1}^j \frac{k}{k+1} b_{j-k} \quad (j \geq 1). \quad (2.4)$$

Use of (2.4) is easily seen to generate the values

$$b_1 = \frac{1}{2}, \quad b_2 = \frac{11}{24}, \quad b_3 = \frac{7}{16}, \quad b_4 = \frac{2447}{5760}, \quad b_5 = \frac{959}{2304}, \quad b_6 = \frac{238043}{580608}, \dots,$$

which are the same coefficients as in (2.1). The representation using a recursive algorithm for the coefficients b_j is more practical for numerical evaluation than the expression in (2.3).

The above result immediately shows that $b_j > 0$ so that (2.2) is an alternating series for positive x . Replacement of x by $1/x$ in (2.1) and (2.2) then enables us to write

$$(1+x)^{1/x} = e \sum_{j=0}^{\infty} (-1)^j b_j x^j \quad (-1 < x \leq 1). \quad (2.5)$$

We now establish a monotonicity property satisfied by the coefficients b_j .

LEMMA 2.1. *The sequence $\{b_j\}_{j=0}^{\infty}$ in (2.5) is monotonically decreasing.*

Proof. By Cauchy's theorem it follows from (2.5) that

$$b_j = \frac{(-1)^j}{2\pi i e} \oint_C (1+t)^{1/t} \frac{dt}{t^{j+1}},$$

where C is a closed loop surrounding $t = 0$ described in the positive sense. Define

$$\Delta_j = b_j - b_{j+1}.$$

Then

$$\Delta_j = \frac{(-1)^j}{2\pi i e} \oint_C (1+t)^{1/t} \left(1 + \frac{1}{t}\right) \frac{dt}{t^{j+1}} = \frac{(-1)^j}{2\pi i e} \oint_C (1+t)^{1+1/t} \frac{dt}{t^{j+2}}.$$

In the t -plane there is a branch cut along $(-\infty, -1]$. Now expand C to be a large circle of radius R that is indented to pass along the upper and lower sides of the branch cut. The contribution from the large circle tends to zero as $R \rightarrow \infty$. Similarly, the contribution round the branch point $t = -1 + \rho e^{i\theta}$, $-\pi \leq \theta \leq \pi$ vanishes as $\rho \rightarrow 0$. Then we have upon putting $t = xe^{\pm\pi i}$ on the upper and lower sides of the branch cut

$$\begin{aligned} \Delta_j &= \frac{1}{2\pi i e} \int_{\infty}^1 (x-1)^{1-1/x} e^{-\pi i/x} \frac{dx}{x^{j+2}} + \frac{1}{2\pi i e} \int_1^{\infty} (x-1)^{1-1/x} e^{\pi i/x} \frac{dx}{x^{j+2}} \\ &= \frac{1}{\pi e} \int_1^{\infty} (x-1)^{1-1/x} \sin(\pi/x) \frac{dx}{x^{j+2}}. \end{aligned} \quad (2.6)$$

Now on the interval $x \in [1, \infty)$ the function $\sin(\pi/x) \geq 0$ so that the integrand in (2.6) is non-negative on $[1, \infty)$. Hence $\Delta_j > 0$ and the sequence $\{b_j\}_{j=0}^\infty$ is monotonically decreasing. This completes the proof.

REMARK 2.1. We thank a referee for providing the literature [1]. It was proved in [1, Lemma 1] that

$$(x+1) \left[e - \left(1 + \frac{1}{x} \right)^x \right] = \frac{e}{2} + \int_0^1 \frac{g(s)}{x+s} ds \quad (x > 0), \quad (2.7)$$

where

$$g(s) = \frac{1}{\pi} s^s (1-s)^{1-s} \sin(\pi s) \quad (0 \leq s \leq 1). \quad (2.8)$$

By (2.7), we here give an integral representation of the coefficients b_j in (2.5), and then use it to prove Lemma 2.1.

Write (2.7) as

$$\left(1 + \frac{1}{x} \right)^x = e - \frac{e}{2(x+1)} - \int_0^1 \frac{g(s)}{(x+1)(x+s)} ds \quad (x > 0). \quad (2.9)$$

Replacing x by $1/t$ in (2.9) yields, for $t > 0$,

$$\begin{aligned} f(t) &:= (1+t)^{1/t} = \frac{e}{2} + \frac{e}{2(t+1)} - \int_0^1 \frac{g(s)}{s} \frac{t^2}{(t+1)(t+\frac{1}{s})} ds \\ &= \frac{e}{2} + \frac{e}{2(t+1)} - \int_0^1 \frac{g(s)}{s} \left\{ 1 + \frac{s}{(1-s)(t+1)} - \frac{1}{s(1-s)(t+\frac{1}{s})} \right\} ds. \end{aligned} \quad (2.10)$$

Clearly,

$$eb_0 = f(0) = e.$$

Differentiating the expression in (2.10), we find that for $n \geq 1$,

$$\frac{(-1)^n f^{(n)}(t)}{n!} = \frac{e}{2(t+1)^{n+1}} - \int_0^1 \frac{g(s)}{s} \left\{ \frac{s}{(1-s)(t+1)^{n+1}} - \frac{1}{s(1-s)(t+\frac{1}{s})^{n+1}} \right\} ds,$$

we then obtain the following integral representation of the coefficients b_j in (2.5):

$$b_n = \frac{(-1)^n f^{(n)}(0)}{n!e} = \frac{1}{2} - \frac{1}{e} \int_0^1 \frac{1-s^{n-1}}{1-s} g(s) ds$$

for $n \geq 1$, and we have

$$\Delta_j = b_j - b_{j+1} = \frac{1}{e} \int_0^1 s^{j-1} g(s) ds > 0 \quad (j \geq 1). \quad (2.11)$$

Noting that $b_0 = 1 > \frac{1}{2} = b_1$ holds, we see that the sequence $\{b_j\}_{j=0}^\infty$ in (2.5) is monotonically decreasing.

In fact, by an elementary change of variable $x = 1/s$ ($0 \leq s \leq 1$), we see that (2.6) \iff (2.11).

From (2.5) and Lemma 2.1 we obtain the following theorem that develops the double inequality (1.3) to produce a general result.

THEOREM 2.1. *For all integers $m \geq 0$,*

$$e \sum_{j=0}^{2m+1} (-1)^j b_j x^j < (1+x)^{1/x} < e \sum_{j=0}^{2m} (-1)^j b_j x^j \quad (0 < x \leq 1), \quad (2.12)$$

or alternatively

$$e \sum_{j=0}^{2m+1} \frac{(-1)^j b_j}{x^j} < \left(1 + \frac{1}{x}\right)^x < e \sum_{j=0}^{2m} \frac{(-1)^j b_j}{x^j} \quad (x \geq 1), \quad (2.13)$$

where the coefficients b_j are given by the recursive relation (2.4).

3. A generalized Carleman-type inequality

THEOREM 3.1. *Let $0 < \lambda_{n+1} \leq \lambda_n$, $\Lambda_n = \sum_{m=1}^n \lambda_m$ ($\Lambda_n \geq 1$), $a_n \geq 0$ ($n \in \mathbb{N}$) and $0 < \sum_{n=1}^{\infty} \lambda_n a_n < \infty$. Then for $0 < p \leq 1$,*

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < \frac{e^p}{p} \sum_{n=1}^{\infty} \left(\sum_{j=0}^{2m} \frac{(-1)^j b_j}{(\Lambda_n/\lambda_n)^j} \right)^p \lambda_n a_n^p \Lambda_n^{p-1} \left(\sum_{k=1}^n \lambda_k (c_k a_k)^p \right)^{(1-p)/p}, \quad (3.1)$$

where b_j is given by (2.4), and

$$c_n^{\lambda_n} = \frac{(\Lambda_{n+1})^{\Lambda_n}}{(\Lambda_n)^{\Lambda_{n-1}}}.$$

Proof. The following inequality:

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} \leq \frac{1}{p} \sum_{m=1}^{\infty} \left(1 + \frac{1}{\Lambda_m/\lambda_m} \right)^{p\Lambda_m/\lambda_m} \lambda_m a_m^p \Lambda_m^{p-1} \left(\sum_{k=1}^m \lambda_k (c_k a_k)^p \right)^{(1-p)/p} \quad (3.2)$$

has been proved in Theorem 2.2 of [11] (see also [21, p. 96]). From (3.2) and the right-hand side of (2.13), we obtain (3.1). The proof is complete.

REMARK 3.1. In Theorem 2.2 of [11], $c_k^{\lambda_k} = \frac{(\Lambda_{k+1})^{\Lambda_k}}{(\Lambda_k)^{\Lambda_{k-1}}}$ should be $c_n^{\lambda_n} = \frac{(\Lambda_{n+1})^{\Lambda_n}}{(\Lambda_n)^{\Lambda_{n-1}}}$; see [11, p. 44, line 3]. Likewise, $c_s^{\lambda_s} = \frac{(\Lambda_{s+1})^{\Lambda_s}}{(\Lambda_s)^{\Lambda_{s-1}}}$ in Theorem 3.1 of [21] should be $c_n^{\lambda_n} = \frac{(\Lambda_{n+1})^{\Lambda_n}}{(\Lambda_n)^{\Lambda_{n-1}}}$; see [21, p. 96, Eq. (9)].

The choice $p = 1$ in (3.1) yields

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < e \sum_{n=1}^{\infty} \left(\sum_{j=0}^{2m} \frac{(-1)^j b_j}{(\Lambda_n/\lambda_n)^j} \right) \lambda_n a_n. \quad (3.3)$$

Taking $\lambda_n \equiv 1$ in (3.3) we obtain

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} \left(\sum_{j=0}^{2m} \frac{(-1)^j b_j}{n^j} \right) a_n. \quad (3.4)$$

When $m = 2$ in (3.4) we recover (1.5).

REFERENCES

- [1] H. ALZER AND C. BERG, *Some classes of completely monotonic functions*, Ann. Acad. Sci. Fenn. Math. **27** (2002), 445–460.
- [2] H. J. BROTHERS AND J. A. KNOX, *New closed-form approximations to the logarithmic constant e* , Math. Intelligencer **20** (1998), 25–29.
- [3] T. CARLEMAN, *Sur les fonctions quasi-analytiques*, Comptes rendus du V^e Congrès des Mathématiciens Scandinaves, Helsingfors (1922), 181–196.
- [4] H.-W. CHEN, *On an infinite series for $(1+1/x)^x$ and its application*, Int. J. Math. Math. Sci. **29** (2002), 675–680.
- [5] C.-P. CHEN, *Generalization of weighted Carleman-type inequality*, East J. Approx. **12** (2006), 63–69.
- [6] C.-P. CHEN, W.-S. CHEUNG AND F. QI, *Note on weighted Carleman type inequality*, Int. J. Math. Math. Sci. **3** (2005), 475–481.
- [7] C.-P. CHEN AND J. CHOI, *Asymptotic formula for $(1+1/x)^x$ based on the partition function*, Amer. Math. Monthly **121** (2014), 338–343.
- [8] C.-P. CHEN, N. ELEZOVIĆ AND L. VUKŠIĆ, *Asymptotic formulae associated with the Wallis power function and digamma function*, J. Classical Anal. **2**, 2 (2013), 151–166.
- [9] C.-P. CHEN AND F. QI, *Generalization of Hardy's inequality*, Proc. Jangjeon Math. Soc. **7** (2004), 57–61.
- [10] A. ČIŽMEŠIJA, J. PEČARIĆ AND L. E. PERSSON, *On strengthened weighted Carleman's inequality*, Bull. Austral. Math. Soc. **68** (2003), 481–490.
- [11] S. S. DRAGOMIR AND Y.-H. KIM, *The strengthened Hardy inequalities and their new generalizations*, Fac. Sci. Math. **20**, 2 (2006), 39–49.
- [12] G. H. HARDY, *Notes on some points in the integral calculus*, Messenger of Math. **54** (1925), 150–156.
- [13] G. H. HARDY, J. E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, 2nd ed., Cambridge Univ. Press, Cambridge, 1952.
- [14] Y. HU, *A strengthened Carleman's inequality*, Commun. Math. Anal. **1** (2006) 115–119.
- [15] M. JOHANSSON, L. E. PERSSON AND A. WEDESTIG, *Carleman's inequality—history, proofs and some new generalizations*, J. Inequal. Pure Appl. Math. **4**, 3 (2003), Art. 53.
http://www.emis.de/journals/JIPAM/images/135_02_JIPAM/135_02_www.pdf.
- [16] S. KAIJSER, L. E. PERSSON AND A. ÖBERG, *On Carleman and Knopp's inequalities*, J. Approx. Theory **117** (2002), 140–151.
- [17] J. A. KNOX AND H. J. BROTHERS, *Novel series-based approximations to e* , College Math. J. **30** (1999), 269–275.
- [18] A. KUFNER AND L. E. PERSSON, *Weighted Inequalities of Hardy Type*, World Scientific, New Jersey/London, Singapore/Hong Kong, 2003.
- [19] J.-L. LI, *Notes on an inequality involving the constant e* , J. Math. Anal. Appl. **250** (2000), 722–725.
- [20] H.-P. LIU AND L. ZHU, *New strengthened Carleman's inequality and Hardy's inequality*, J. Inequal. Appl. J. Inequal. Appl. **2007**, Art. ID 84104, 7 pp.
<http://link.springer.com/article/10.1155/2007/84104/fulltext.html>.

- [21] Z. LÜ, Y. GAO AND Y. WEI, *Note on the Carleman's inequality and Hardy's inequality*, Comput. Math. Appl. **59** (2010), 94–97.
- [22] C. MORTICI AND Y. HU, *On some convergences to the constant e and improvements of Carleman's inequality*, Carpathian J. Math. **31** (2015), 249–254.
- [23] C. MORTICI AND X.-J. JANG, *Estimates of $(1+x)^{1/x}$ Involved in Carleman's Inequality and Keller's Limit*, Filomat **29**, 7 (2015), 1535–1539.
- [24] J. PEČARIĆ AND K. B. STOLARSKY, *Carleman's inequality: history and new generalizations*, Aequationes Math. **61** (2001), 49–62.
- [25] G. PÓLYA, *Proof of an inequality*, Proc. London Math. Soc. **24** (1926), 57.
- [26] Z. XIE AND Y. ZHONG, *A best approximation for constant e and an improvement to Hardy's inequality*, J. Math. Anal. Appl. **252** (2000), 994–998.
- [27] P. YAN AND G.-Z. SUN, *A strengthened Carleman's inequality*, J. Math. Anal. Appl. **240** (1999), 290–293.
- [28] B.-C. YANG AND L. DEBNATH, *Some inequalities involving the constant e , and an application to Carleman's inequality*, J. Math. Anal. Appl. **223** (1998), 347–353.
- [29] B.-C. YANG, *On Hardy's inequality*, J. Math. Anal. Appl. **234** (1999), 717–722.
- [30] X.-J. YANG, *On Carleman's inequality*, J. Math. Anal. Appl. **253** (2001), 691–694.
- [31] X.-J. YANG, *Approximations for constant e and their applications*, J. Math. Anal. Appl. **262** (2001), 651–659.

(Received November 12, 2016)

Chao-Ping Chen
School of Mathematics and Informatics
Henan Polytechnic University
Jiaozuo City 454000, Henan Province, China
e-mail: chenchao ping@sohu.com

Richard B. Paris
Division of Computing and Mathematics
University of Abertay
Dundee, DD1 1HG, UK
e-mail: R.Paris@abertay.ac.uk

